

## PERSPECTIVE

Q: 1, 2, 3, 4, 5

# Reduced development of COVID-19 in children reveals molecular checkpoints gating pathogenesis illuminating potential therapeutics

Jonathan Baruch Steinman<sup>a</sup>, Fok Moon Lum<sup>b,c</sup>, Peggy Pui-Kay Ho<sup>b,c</sup>, Naftali Kaminski<sup>d</sup>, and Lawrence Steinman<sup>b,c,1</sup>

Edited by Robert L. Coffman, University of California, Santa Cruz, Portola Valley, CA, and approved August 14, 2020 (received for review July 8, 2020)

**The reduced development of COVID-19 for children compared to adults provides some tantalizing clues on the pathogenesis and transmissibility of this pandemic virus. First, ACE2, the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) Receptor, is reduced in the respiratory tract in children. Second, coronavirus associated with common colds in children may offer some protection, due to cross-reactive humoral immunity and T cell immunity between common coronaviruses and SARS-CoV-2. Third, T helper 2 immune responses are protective in children. Fourth, surprisingly, eosinophilia, associated with T helper 2, may be protective. Fifth, children generally produce lower levels of inflammatory cytokines. Finally, the influence of the downturn in the global economy, the impact of living in quarters among families who are the most at risk, and factors including the openings of some schools, are considered. Those most disadvantaged socioeconomically may suffer disproportionately with COVID-19.**

COVID-19 | children | SARS-CoV-2 | ACE2

**Q:7** Our children, often a source of hope and optimism, have provided one of the few encouraging notes, a “silver lining” perhaps, in the dismal landscape of the current pandemic. Children have thus far been relatively protected from developing severe COVID-19 pneumonia. Understanding the basis for why children have generally done better than adults provides important insights into the pathogenesis of COVID-19, which may guide our understanding of susceptibility to infection and may provide further clues for therapeutics.

**Q:9** In the United States, through April 2, 2020, two percent of the reported COVID-19 cases were in children less than 18 y of age, according to the Centers for Disease Control and Prevention (CDC) (1). However, as of August 6, 2020, the American Academy of Pediatrics reported 9.1% of all cases of COVID-19 were children in states reporting cases by age. This figure represents 380,000 children in the United States, who have tested positive for COVID-19 since the onset of the pandemic

(2). This increase over 4 mo is noteworthy. Whatever the underlying reasons for this increase may be, this trend serves to remind us that we are in an early phase of understanding the impact of this virus on children.

The CDC reported through the first week of August 2020 that, since March 1, 2020, there have been 576 pediatric COVID-19-associated hospitalizations, reported through the CDC’s COVID-NET (3). Studies from China, Italy, Spain, and North America all indicate that children are less frequently hospitalized with COVID-19 than adults (1–7). Evidence from the North American study (6) shows that many of the children who did require inpatient care due to acute COVID-19 had preexisting medical problems, including conditions that required long-term dependence on technologic support for their underlying diseases, as well as comorbidities that included obesity, immune suppression, and cancer (6). Hispanic or Latino (Hispanic)

<sup>a</sup>Department of Pediatrics, Columbia University, New York, NY 10032; <sup>b</sup>Department of Pediatrics, Stanford University, Stanford CA 94305; <sup>c</sup>Department of Neurology & Neurological Sciences, Stanford University, Stanford, CA 94305; and <sup>d</sup>Pulmonary, Critical Care and Sleep Medicine, Yale University School of Medicine, New Haven, CT 06520

Author contributions: J.B.S., F.M.L., P.P.-K.H., N.K., and L.S. analyzed data and wrote the paper.


Competing interest statement: N.K. has consulted with Biogen Idec, Boehringer Ingelheim, Third Rock, Pliant, Samumed, NuMedii, Indaloo, Theravance, LifeMax, Three Lake Partners, RohBar, and reports a grant from VeracYTE. N.K. has patents on New Therapies in Pulmonary Fibrosis and Peripheral Blood Gene Expression Biomarkers. All these consultations are outside of this work. L.S. has consulted with Roche, Novartis, Tolerion, Atreca, TG Therapeutics, and Atara Biopharma. All these consultations are outside of the scope of this work.

This article is a PNAS Direct Submission.

This open access article is distributed under [Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 \(CC BY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/).

<sup>1</sup>To whom correspondence may be addressed. Email: steinman@stanford.edu.

## Five Clues Why Children Have Reduced Susceptibility to COVID-19



**Fig. 1. (1) Coronavirus associated with common colds in children may offer some protection due to cross-reactive T cell immunity and cross-reactive antibody immunity between common coronaviruses and SARS-CoV-2, and due to reduced ACE2 in nasal mucosa of children. (2) Reduced TMPRSS2 in children in type I alveolar cells (AEI). Reduced ACE2 in children in type II alveolar cells (AEII). (3) Protective Th2 immunity in children. (4) Surprising protection from eosinophilia driven by Th2 cytokines including IL-4, IL-5 and IL-13 (for example childhood asthma). (5) Children produce lower levels of inflammatory cytokines, IL-6 production increases with age.**

children and non-Hispanic Black children were hospitalized at rates approximately eightfold and fivefold higher than non-Hispanic White children (3).

Although relatively rare, a multisystem inflammatory syndrome in children (abbreviated MIS-C) has been reported from pediatric departments in hospitals. A large majority of children with MIS-C were positive, with PCR or with serologic testing, for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (7). It is unclear, at this point, whether the pathogenesis reflects active infection, given

the variable presence of a positive PCR, but, more likely, the syndrome reflects predominately a postinfectious immune response. The percentage of Black and Hispanic children with MIS-C was higher than seen in the US population (8, 9) Thankfully, mortality related to this complication of SARS-CoV-2 has remained low (~2%) (8).

Exploring why the pediatric population is generally far less likely to develop COVID-19, even though their rate of infection is similar to adults (10), may offer productive clues, enabling strategies for targeting key pathogenic mechanisms in our efforts to contain and

249 eradicate transmission of the virus. Here we review some of these  
250 intriguing areas that may help illuminate why children might be  
251 more resistant to serious outcomes following exposure to SARS-  
252 CoV-2.

### 253 ACE2 Expression Is Lower in the Respiratory Tracts of 254 Children

255 Both the first SARS virus, SARS-CoV-1 discovered in 2003, and the  
256 SARS-CoV-2 virus bind to angiotensin-converting enzyme 2 (ACE2)  
257 (11, 12). The renin angiotensin system, originally known for its  
258 critical role in blood pressure and hypertension, is also a critical  
259 trigger of inflammation in various organ systems. ACE converts the  
260 10-amino acid angiotensin I peptide into the 8-amino acid peptide,  
261 known as angiotensin II. ACE2 then converts angiotensin II to the 7-  
262 amino acid peptide known as angiotensin<sub>1-7</sub>. Angiotensin-II, when  
263 binding to the angiotensin type (AT) 1 receptor, is proinflammatory  
264 in multiple organ systems including the lungs, the heart, the kidneys  
265 (13, 14), and the brain (15). In contrast, angiotensin<sub>1-7</sub> is anti-  
266 inflammatory when it binds to the Mas receptor (14).

267 Coronaviruses are enveloped single-stranded RNA viruses that  
268 cause enteric and respiratory tract infections in mammals and  
269 birds that can range from mild to severe (16). After the spike  
270 protein of SARS-CoV-2 binds to cells in the respiratory tract via  
271 ACE2 (11–13, 17, 18), the virus enters the cell via proteolytic  
272 cleavage involving two proteases, the transmembrane protease  
273 serine 2 (TMPRSS2) and cathepsin L (CTSL). In recent studies (17,  
274 18) researchers analyzed the expression of RNA in cells of the  
275 respiratory tract, including nasal, airway, and lung parenchyma.  
276 Both of these studies showed that expression of ACE2 increases  
277 with age in the respiratory tract.

278 Wang et al. (17) analyzed gene expression in healthy lung tissue  
279 from three different age groups ranging from approximately age  
280 30 gestational weeks, approximately age 3 y, and approximately  
281 age 30 y. They analyzed gene expression with two platforms at  
282 single nuclear (sn) resolution, using snRNA sequencing (snRNA-seq)  
283 and sn assay for transposase-accessible chromatin (snATAC-seq),  
284 enabling the researchers to analyze specific genes and neighboring  
285 (cis-) regions that control cell-type expression of gene modules in-  
286 volved in processes like inflammation.

287 The respiratory tract has diverse cell types including alveolar  
288 epithelial type 1 (AEI) cells, that are responsible for the air–blood  
289 barrier in lung, and alveolar epithelial type 2 (AEII) cells that are  
290 critical for secreting pulmonary surfactant (17, 18). The cellular  
291 distribution of ACE2 and TMPRSS2 was distinct from CTSL. While  
292 ACE2 and TMPRSS2 were primarily in AEI, AEII, and airway cells  
293 such as club, ciliated, and basal cells, CTSL expression was found in  
294 epithelial cells, mesenchymal cells, endothelial cells, and macro-  
295 phages. It is the alveolar cells that are the anatomic site of acute  
296 respiratory distress syndrome.

297 Wang et al. (17) noted “an increase in the proportion of alveolar  
298 epithelial cells expressing ACE2 and TMPRSS2 in adult compared  
299 to young lungs.” The investigators observed marked differences,  
300 particularly in AE cells (17), with the percentage of ACE2<sup>+</sup> AEII cells  
301 increasing with age from the samples from those age 3 y upward to  
302 the samples from those age 30 y. Similar findings were seen for the  
303 TMPRSS2<sup>+</sup> cells.

304 Analysis of the regulatory elements related to TMPRSS2<sup>+</sup> cells  
305 revealed modules related to activation of an immune response in  
306 the lungs with a correlation related to increasing age. Noteworthy  
307 immune factors included interferon (IFN) regulatory molecules for  
308 both type 1 and type 2 IFNs, and signal transducer and activator  
309 of transcription molecules involved in activating inflammatory

310 cytokine molecular pathways. Expression of TMPRSS2 was highest  
311 in ciliated cells and AE1. Expression of TMPRSS2 increased with  
312 age in mice and humans (19). For SARS-CoV-1, TMPRSS2 was  
313 shown to promote viral spread, with its protease activity serving to  
314 reduce viral recognition by neutralizing antibodies. Whether  
315 TMPRSS2 plays a similar role in the spread of SARS-CoV-2 is a  
316 subject of current investigation (19).

317 Muus et al. (18) analyzed gene expression in more than 4  
318 million cells derived from 107 single-cell RNA expression studies,  
319 including 22 studies focusing on datasets from the lung and air-  
320 ways in both adults and children. In this large metaanalysis of  
321 multiple RNA expression studies, the researchers compiled data  
322 on cells from “nasal, airway, and lung parenchyma samples from  
323 164 donors spanning fetal, childhood, adult, and elderly age  
324 groups.” They found ACE2, TMPRSS2, and CTSL coexpressed at  
325 low levels in alveolar cells of children compared to adults. The  
326 researchers were able to confirm these results directly with triple  
327 fluorescence in situ hybridization in specific cell types. They  
328 concluded that there was “particularly low” ACE2 in the pediatric  
329 samples (18).

330 Although the number of samples of young children was limited  
331 in this study (18), the researchers found that “strikingly,  
332 ACE2 expression is very low in normal lungs of newborns, in the  
333 sample of the individual of 3 mo, and in the lungs of the 3-y-olds.”  
334 These data were supported on a second platform known as  
335 scTHS-Seq, single-cell chromatin accessibility by transposome  
336 hypersensitive site sequencing, from human pediatric samples  
337 with no lung disease, collected at day 1 of life, at 14 mo, at 3 y,  
338 and at 9 y. They found “no signal was present at birth, it was low in  
339 the 9-y-old and 3-y-old sample, and higher in the 14 mo-old.” The  
340 researchers concluded that “these patterns may be important in  
341 understanding why children are more resistant to COVID-19” (18).

342 There was an association of cells expressing both ACE2 and  
343 TMPRSS2 that were more frequent with increasing age (18). Cells  
344 expressing both TMPRSS2 and ACE2 in children are quite rare  
345 (18). These double-positive cells were particularly notable for  
346 expression of associated modules mediating viral and immune  
347 responses. These double-positive cells were the likely locus where  
348 an exuberant inflammatory response emanates. In these double-  
349 positive cells in lung epithelium, expression of interleukin 6 (IL-6)  
350 and IL-6R was increased. In so-called cytokine storms, where there  
351 is overactivation of the immune system in the alveoli, activation of  
352 these associated gene modules may underlie the pathophysiologic  
353 response (18). Monoclonal antibodies targeting inflamma-  
354 tory cytokines including IL-6 and its receptor, as well as tumor  
355 necrosis factor (TNF), are in therapeutic trials for cytokine storm  
356 in COVID-19.

357 A reasonable conjecture might be that, if ACE2 and/or  
358 TMPRSS2 expression is diminished in children, then viral infection  
359 of respiratory cells by SARS-CoV-2 might be less likely at any given  
360 viral load, and, additionally, there might be reduced expression of  
361 associated inflammatory modules. It is known that, in healthy  
362 children, there is an age-related increase in the production of IFN- $\gamma$   
363 and TNF- $\alpha$ . For IL-6, there was a trend for lower IL-6 concentrations  
364 in children below 2 y of age compared to older children (20). Thus,  
365 children have a lowered propensity to transcribe inflammatory cy-  
366 tokines in lung cells, and this may reduce the chance for an ex-  
367 plosive immune reaction within the lungs resulting in the so-called  
368 “cytokine storm.”



## Impact of Frequent Respiratory Infections in Childhood and Virologic and Immunologic Interference

Numerous studies (21–23) show that children younger than 2 y of age have five or more respiratory infections per year, and spend a median of 44 d with mild upper respiratory illnesses (21). The activation of adaptive immunity to common coronaviruses as well as the activation of the innate immune system in the respiratory tract may indeed provide some protection from microbial infection, including SARS-CoV-2. The frequent respiratory infections in children may provide further clues to why children are resistant. One line of reasoning is based on the phenomenon of viral interference. A second line of reasoning is based on the concept of immune interference.

**Viral Interference.** Often, children are infected by more than one viral agent (21, 22). Viral interference is a well-known phenomenon where one virus interferes with the replication of a second virus (23). There is some evidence for coinfections in COVID-19 patients, including coinfection with other coronaviruses (24, 25). Kim et al. (24) reported on both hospitalized and nonhospitalized cases from Northern California, including children. They found more than 20% were positive for a second viral infection, including infection with other coronaviruses. In contrast, Nowak et al. (25) reported concurrent viral infection in only 3% of 1,204 SARS-CoV-2–positive patients from the New York metropolitan area who were also tested with a respiratory virus panel or a test for influenza and respiratory syncytial virus (RSV). In comparison, of 7,418 patients who tested negative for SARS-CoV-2, 845 were tested with the same multiplex panels, and 13% were positive for at least one non-SARS-CoV-2 respiratory viral pathogen (25). The increase in at least one non-SARS-CoV-2 viral pathogen in those who tested negative for SARS-CoV-2 may be due, in part, to viral interference.

An intriguing potential mechanism of resistance in children is that common coronaviruses associated with mild illnesses like colds and more severe illness like croup and bronchiolitis are associated with decreased expression of ACE2. For example, human coronavirus (HCoV) NL63, associated with common colds and croup, induces a down-regulation of ACE2 (26). A reduction in the viral receptor for SARS-CoV-2 therefore might help explain why children who carry such viruses in their nose and upper portions of their respiratory tracts are hospitalized less frequently than adults. The first site of encounter in the respiratory tract for the SARS-CoV-2 is in the nose. Investigators studying a cohort of 305 patients aged 4 y to 60 y found that “older children (10 y to 17 y old;  $n = 185$ ), young adults (18 y to 24 y old;  $n = 46$ ), and adults ( $\geq 25$  y old;  $n = 29$ ) all had higher expression of ACE2 in the nasal epithelium compared with younger children (4 y to 9 y old;  $n = 45$ )” (27).

Despite diminished expression of ACE2 and/or TMPRSS2 in children, “symptomatic infants have higher nasopharyngeal SARS-CoV-2 viral loads at presentation but develop less severe disease than older children and adolescents” (28). Moreover, children less than 5 y old “with mild to moderate COVID-19 have high amounts of SARS-CoV-2 viral RNA in their nasopharynx compared with older children and adults” (29).

Taken together, these data imply that, although the viral load in the nasopharynx may be higher in symptomatic children, overall, children remain markedly more resistant to viral infection of the lower respiratory tract leading to COVID-19. All this raises the likelihood that, although children are, fortunately, less susceptible to viral infection of the lungs, they can still serve as good disseminators of the SARS-CoV-2 virus that lurks in their nasal mucosa.

Detailed contact tracing from the Korean CDC revealed that household contacts of COVID-19–positive children ages 10 y to 19 y were the most likely to become infected with the SARS-CoV-2 virus compared to household contacts of people of all other ages. In contrast, household contacts of positive children aged 0 y to 9 y were the least likely to become infected. Taken together, these data need further substantiation, and might have implications for spread of infection at home, in daycare, and in schools (30). These data all emphasize that there is much to learn and to consider on the spread from children in various environments. It will also be important to stratify children by age, as infants, young children, and adolescents have different propensities for communicating viral infection.

**Immunological Interference.** IFNs, first described in 1957 by Isaacs and Lindemann, bind to three distinct types of IFN receptors (31). Mapping of immune response modules in the respiratory tract shows that gene modules triggered by both type 1 and type 2 IFN responses are prominent (17, 18, 32, 33). The type 1 IFN response is vital for viral killing. Type 1 IFN, however, stimulates expression of the ACE2 receptor for the SARS-CoV-2 virus (33). Thus, the virus drives an increase in type 1 IFN expression, which then enhances expression of its receptor in the airway (33). In contrast, coronaviruses that frequently infect children with common colds down-regulate ACE2 as described above (26). Thus, children may benefit from a virtuous cycle with decreased ACE2 leading to less induction of the IFN response, which, in turn, further attenuates ACE2 expression. In contrast, adults suffer from a vicious cycle in which increased ACE2 expression drives a more robust IFN response.

The adaptive immune response to common coronavirus infections in children could provide some protection to SARS-CoV-2 since they share considerable degrees of homology with coronaviruses associated with the common cold. For example, spike proteins of the common HCoV share 30% amino acid identity with these viruses when one performs BLAST searches comparing these viruses. A detailed mapping of known T and B cell epitopes on SARS-CoV-2 indicates that adaptive immune reactivity at the T cell and antibody level targets not just the spike region but also other viral proteins (34).

A study in adult donors, age currently greater than 20 y, tested whether there was detectable immunity to SARS-CoV-2 and to common cold viruses attributed to coronavirus strains HCoV-OC43, HCoV-HKU1, HCoV-NL63, and HCoV-229E. Donors were recruited between 2015 and 2018, obviating exposure to SARS-CoV-2. Unexposed donors had T cell immunity to both spike and nonspike proteins in SARS-CoV-2 (34). The immunity may have emanated from shared regions on the common cold virus and SARS-CoV-2. Investigators tested immunity to one betacoronavirus HCoV-OC43 and to one alphacoronavirus NL63, and showed that these unexposed donors,  $n = 11$ , all had IgG to these to the receptor-binding domain of these common cold viruses. These nonexposed donors also had vigorous T cell responses to both spike and nonspike proteins in SARS-CoV-2 (35). Additional studies have been reported demonstrating widespread immunity in healthy individuals to cross-reactive regions of SARS-CoV2 that share peptide sequence homology with endemic coronaviruses that cause common colds like HCoV-OC43, HCoV-229E, HCoV-NL63, and HCoV-HKU1 (36, 37). Humoral, antibody immunity to SARS-CoV2 is widely detected in individuals, including children, who were not exposed to SARS-CoV2 (38). Immunity to earlier exposures to both alpha and beta coronaviruses may thus

engender protective humoral and cellular immunity for children who are heavily exposed to these common cold viruses.

A publication (39) from the Department of Defense examining the effect of the 2017–2018 seasonal influenza vaccine on respiratory infections produced some intriguing results. Investigators showed that the seasonal influenza immunization protected against influenza, as it was intended to do, and protected against some other respiratory viruses. However, they noted a small but statistically significant increase in individuals testing positive to metapneumovirus and coronaviruses. If future influenza vaccines, for example the 2020–2021 seasonal influenza vaccine, also provide increased occurrence of common coronaviruses, this phenomenon may actually afford some protection to SARS-CoV-2. Recent studies show that those with immunity to common coronaviruses do have adaptive immunity to SARS-CoV-2 via the mechanism of cross-reactive immunity (35–37).

There is some precedent for this phenomenon. Some immunizations induce protection against other infections outside of the intended target of the vaccine itself (40). A study from the Mayo Clinic indicated that “polio, Hemophilus influenzae type-B (HIB), measles-mumps-rubella (MMR), varicella, pneumococcal conjugate (PCV13), geriatric flu, and hepatitis A/hepatitis B (HepA-HepB) vaccines administered in the past 1, 2, and 5 y are associated with decreased SARS-CoV-2 infection rates” (41).

Some antiviral vaccines like the MMR vaccine contain components that have structural similarities with SARS-CoV-2 (42, 43). There is a 29% amino acid sequence homology between the ADP ribose-1-phosphatase domains of SARS-CoV-2 and rubella virus, including surface-exposed conserved residues shared between SARS-CoV-2 and the attenuated rubella virus in MMR (42). Patients with COVID-19 infection had raised levels of rubella IgG, but did not have increased rubella IgM, nor did they have increased levels of antibody to varicella zoster. The investigators interpret these results as indicative of a cross-reactive recall antibody response, common to regions shared between rubella and SARS-CoV-2, that may modulate the course of disease in COVID-19 (42). Whether or not such an immune response is protective or whether such an immune response might potentially enhance disease are outcomes under investigation (42).

### A Surprising Potential Protective Benefit of Th2 Immunity in Children

There are three major arms of the immune response characteristics of human T helper cells, named Th1, Th2, and Th17 (33). The Th1 arm described above is mediated by gamma IFN. The Th2 arm is associated with allergic disease, and is mediated by IL-4, IL-5, and IL-13 (44). Sajuthi et al. (33) reported on gene expression studies on 695 children with asthma and healthy controls from the Genes-Environment & Admixture in Latino Americans study, an ongoing case-control study of asthma in Latino children and adolescents. They found that TMPRSS2 is part of a mucus secretory network, driven by Th2 inflammation via the actions of IL-13 (33). They found that Th2 responses driven by IL-4, IL-5, and IL-13 “dramatically” reduced ACE2 in the respiratory tract and are associated with better clinical outcomes with COVID-19, while the type 1 IFN response to respiratory viruses increased ACE2 expression (33).

Th2 cytokines drive an increase of a cell type called the eosinophil in the blood and tissues. Eosinophilia is a hallmark of Th2 inflammation in the airways, most notably in asthma (33, 45). Sajuthi et al. (33) concluded that, at least “provisionally... T[h]2 inflammation may predispose individuals to experience better COVID-19 outcomes through a decrease in airway levels of ACE2 that

override any countervailing effect from increased expression of TMPRSS2.” It is indeed surprising that the Th2 immune type associated with allergic disease including asthma, and with eosinophilia, provides some protection to COVID-19 in children. These findings from Sajuthi et al. (33) on 695 children and adolescents may help explain why low levels of eosinophilia were seen in fatalities in the elderly. Du et al. (45) reported that, in a study of 85 fatal COVID-19 adult subjects, 81.2% exhibited very low levels of blood eosinophils. A connection between knowledge gained in studying children (33) may help explain surprising outcomes in the elderly (45). This is one of the unexpected benefits of investigations on the “extremes of outcome” based on comparing children with the elderly populations at highest risk.

Another independent verification of the possible protective role of Th2 immunity was seen in a study on MIS-C. The very low IgE levels seen in individuals with MIS-C indicate that they may have lacked an adequate Th2 response to attenuate the increased inflammation associated with this hyperinflammatory complication in children (46).

One important caveat about the potential protective effect of eosinophilia comes from studies done 50 y ago in making a vaccine against RSV. “In 1967, infants and toddlers immunized with a formalin-inactivated vaccine against RSV experienced an enhanced form of RSV disease characterized by high fever, bronchopneumonia, and wheezing when they became infected with wild-type virus in the community. Hospitalizations were frequent, and two immunized toddlers died upon infection with wild-type RSV. The enhanced disease was initially characterized as a ‘peribronchiolar monocytic infiltration with some excess in eosinophils’” (47). As new SARS-CoV2 vaccines are tested, the potential appearance of Th2 immunity and eosinophilia must be scrutinized through a lens of caution. Although Th2 responses appear to be associated with some degree of protection to COVID-19, based on the experience with development of a novel vaccine to RSV 50 y ago, the dreaded development of immune enhancement, rather than immunization, must be assessed.

### Unanswered Questions and Lessons to Learn

Although children have milder outcomes, thus far in the first few months of the COVID-19 pandemic, much remains unknown (1–10). The list of questions is considerable. We do not know, at this point, whether children who are asymptomatic and who are major carriers of virus can spread disease. When children leave a “shelter-in-place” environment and return to school, how will this impact more-vulnerable populations? Recent reports of outbreaks at summer camps and schools raise serious concerns (48).

The emergence of MIS-C is particularly worrisome, and it is unclear whether there will be long-term sequelae of SARS-CoV-2 infection in patients who were asymptomatic or mildly symptomatic, in those who developed severe COVID-19, or in those who subsequently developed MIS-C (5–9, 49). Other pandemics have been associated with postinfectious phenomena including postencephalitic Parkinson’s in the 1918 influenza pandemic (50). However, the simultaneous emergence of MIS-C in this current pandemic may allow elucidation of the pathophysiological underpinnings of this COVID-19 syndrome with systemic manifestations. Studies on MIS-C might provide insightful comparisons with the Kawasaki syndrome that has some clinical similarities. It is important to remember that intense inflammation activates the coagulation cascade, driving it in a procoagulant direction. Coagulation disorders and thrombosis are seen in MIS-C (51, 52). Autopsy studies on adults have revealed intense thrombotic

microangiopathy with accompanying abnormalities in the coagulation cascade (52). The nexus between the coagulation cascade and the inflammatory response perhaps deserves increased attention for the possible therapeutic modalities that emerge from such a perspective (53, 54). Increased research emphasis on the connection between coagulation and inflammation is recommended.

The types of comorbidity underlying those at risk for COVID-19 accentuate some major differences between adults and children. Although children, like adults, with COVID-19 had a high incidence of comorbid conditions, there were significant differences in the types of comorbidity. The most common comorbidity in children, 40%, was those who were dependent on technological support. These children on technological support had congenital conditions, including developmental delay and genetic anomalies (4). In contrast, the comorbid conditions in adults are often considered acquired, including hypertension, pulmonary disease from smoking, and obesity. One comorbidity common to adults and children is obesity. Obesity was considered a risk factor in 48% of adults, while a lower percentage, 20.5%, of children with COVID-19 were obese (6, 55).

We should remember that, although there is widespread relief that children are hospitalized far less frequently than adults due to COVID-19 infection, we do not know how the carrier status of children affects their caregivers. We again have some relief that studies on pregnant women who test positive for the SARS-CoV-2 virus show that they and their offspring have, so far, done well. There appears to be negligible intrauterine transmission of disease, although larger studies are necessary to exclude this dreaded possibility (56).

While it seems that children are spared the most severe disease manifestations of COVID-19, they will undoubtedly be seriously impacted in many domains, beyond the direct effects of the pathogen. These potential negative impacts include how the pandemic and social distancing from other children affect psychological health, how it impacts education, and how it affects body

weight and childhood obesity when getting sufficient exercise is more difficult during shelter-in-place and safe-distancing restrictions. One wonders how closure of playgrounds may further impact the obesity epidemic in children. Active programs to reduce childhood obesity have been designed so that this comorbidity does not further exacerbate the risk of COVID-19 in children (57). How the pandemic has changed routine medical and dental care, including immunizations to other preventable illnesses, needs to be examined.

The economic crisis associated with the pandemic has its own toll on children, as it is well known that deterioration in social determinants of health has significant adverse effects on children (<https://www.cdc.gov/socialdeterminants/>). Finally, sheltering in place and social distancing may even impact the frequency of infections typically seen in children (21–23). We shall learn, for example, whether these frequent respiratory infections had provided some benefit to children as they form their immune repertoires and gain immune memory.

Dedicating adequate resources to understand the molecular and immune underpinnings of COVID-19 in children may answer many of these questions and potentially allow development of novel therapeutic strategies for enhancement of host resistance to viral infections. We might consider that we may learn as much or more about this virus, from studying the relative resistance to COVID-19 in children, as we will learn from studying the vastly increased level of susceptibility in adults. There is still much to learn.

**Data Availability.** All study data are included in the article.

### Acknowledgments

We were inspired by the efforts of those working in the front line and by the efforts of those behind the front lines, who are supporting one another during this pandemic. We thank Jason Ooi for his creativity, inspiration, and persistence in working with us on the figures. The work was funded by the NIH. L.S. is the Zimmermann Chair of Pediatrics and Neurological Sciences at Stanford. N.K. is the Boehringer Ingelheim Professor of Critical Care and Sleep Medicine at Yale.

- Centers for Disease Control and Prevention, Coronavirus disease 2019 (COVID-19): Caring for children. [https://www.cdc.gov/coronavirus/2019-ncov/hcp/pediatric-hcp.html#anchor\\_1589580133375](https://www.cdc.gov/coronavirus/2019-ncov/hcp/pediatric-hcp.html#anchor_1589580133375). Accessed 11 August 2020.
- American Academy of Pediatrics, Children and COVID-19: State-level data report. <https://services.aap.org/en/pages/2019-novel-coronavirus-covid-19-infections/children-and-covid-19-state-level-data-report/>. Accessed 11 August 2020.
- L. Kim et al., Hospitalization rates and characteristics of children aged <18 years hospitalized with laboratory-confirmed COVID-19—COVID-NET, 14 states, March 1–July 25, 2020. *Morbidity and Mortality Weekly Report*. Accessed 7 August 2020.
- N. Parri, M. Lenge, D. Buonsenso; Coronavirus Infection in Pediatric Emergency Departments (CONFIDENCE) Research Group, Children with covid-19 in pediatric emergency departments in Italy. *N. Engl. J. Med.* **383**, 187–190 (2020).
- S. Richardson et al.; and the Northwell COVID-19 Research Consortium, Presenting characteristics, comorbidities, and outcomes among 5700 patients hospitalized with COVID-19 in the New York city area. *JAMA* **323**, 2052–2059 (2020).
- L. S. Shekardemian et al., Characteristics and outcomes of children with coronavirus disease 2019 (COVID-19) infection admitted to US and Canadian pediatric intensive care units. *JAMA Pediatr.*, 10.1001/jamapediatrics.2020.1948 (2020).
- E. W. Cheung et al., Multisystem inflammatory syndrome related to COVID-19 in previously healthy children and adolescents in New York City. *JAMA*, 10.1001/jama.2020.10374 (2020).
- E. M. Dufort et al.; New York State and Centers for Disease Control and Prevention Multisystem Inflammatory Syndrome in Children Investigation Team, Multisystem inflammatory syndrome in children in New York state. *N. Engl. J. Med.* **383**, 347–358 (2020).
- L. R. Feldstein et al.; Overcoming COVID-19 Investigators and the CDC COVID-19 Response Team, Multisystem inflammatory syndrome in U.S. children and adolescents. *N. Engl. J. Med.* **383**, 334–346 (2020).
- Q. Bi et al., Epidemiology and transmission of COVID-19 in 391 cases and 1286 of their close contacts in Shenzhen, China: A retrospective cohort study. *Lancet Infect. Dis.* **20**, 911–919 (2020).
- W. Li et al., Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. *Nature* **426**, 450–454 (2003).
- A. C. Walls et al., Structure, function, and antigenicity of the SARS-CoV-2 spike glycoprotein. *Cell* **181**, 281–292.e6 (2020).
- Y. Imai, K. Kuba, J. M. Penninger, The discovery of angiotensin-converting enzyme 2 and its role in acute lung injury in mice. *Exp. Physiol.* **93**, 543–548 (2008).
- P. Verdecchia, C. Cavallini, A. Spanevello, F. Angeli, The pivotal link between ACE2 deficiency and SARS-CoV-2 infection. *Eur. J. Intern. Med.* **76**, 14–20 (2020).
- T. V. Lanz et al., Angiotensin II sustains brain inflammation in mice via TGF-beta. *J. Clin. Invest.* **120**, 2782–2794 (2010).
- A. R. Fehr, S. Perlman, Coronaviruses: An overview of their replication and pathogenesis. *Methods Mol. Biol.* **1282**, 1–23 (2015).
- A. Wang et al., Single nucleus multiomic profiling reveals age-dynamic regulation of host genes associated with SARS-CoV-2 infection. [bioRxiv:10.1101/2020.04.12.037580](https://doi.org/10.1101/2020.04.12.037580) (14 April 2020).



745 18 C. Muus *et al.*, Integrated analyses of single-cell atlases reveal age, gender, and smoking status associations with cell type-specific expression of mediators of SARS-CoV-2 viral entry and highlights inflammatory programs in putative target cells. *bioRxiv*:10.1101/2020.04.19.049254 (20 April 2020). 807

746 19 B. A. Schuler *et al.*; Vanderbilt COVID-19 Consortium Cohort; HCA Lung Biological Network, Age-related expression of SARS-CoV-2 priming protease TMPRSS2 in the developing lung. *bioRxiv*:2020.05.22.111187 (23 May 2020). 808

747 20 M. L. Decker, M. P. Grobusch, N. Ritz, Influence of age and other factors on cytokine expression profiles in healthy children—A systematic review. *Front Pediatr.* **5**, 809

748 255 (2017). 810

749 21 L. Toivonen *et al.*, Burden of recurrent respiratory tract infections in children: A prospective cohort study. *Pediatr. Infect. Dis. J.* **35**, e362–e369 (2016). 811

750 22 T. Chonmaitree *et al.*, Viral upper respiratory tract infection and otitis media complication in young children. *Clin. Infect. Dis.* **46**, 815–823 (2008). 812

751 23 N. Kumar, S. Sharma, S. Barua, B. N. Tripathi, B. T. Rouse, Virological and immunological outcomes of coinfections. *Clin. Microbiol. Rev.* **31**, e00111-17 (2018). 813

752 24 D. Kim, J. Quinn, B. Pinsky, N. H. Shah, I. Brown, Rates of co-infection between SARS-CoV-2 and other respiratory pathogens. *JAMA* **323**, 2085–2086 (2020). 814

753 25 M. D. Nowak, E. M. Sordillo, M. R. Gitman, A. E. Paniz Mondolfi, Co-infection in SARS-CoV-2 infected patients: Where are influenza virus and rhinovirus/enterovirus? *J. Med. Virol.*, 10.1002/jmv.25953 (2020). 815

754 26 R. Dijkman *et al.*, Replication-dependent downregulation of cellular angiotensin-converting enzyme 2 protein expression by human coronavirus NL63. *J. Gen. Virol.* **93**, 1924–1929 (2012). 816

755 27 S. Bunyavanich, A. Do, A. Vicencio, Nasal gene expression of angiotensin-converting enzyme 2 in children and adults. *JAMA* **323**, 2427–2429 (2020). 817

756 28 P. Zachariah *et al.*, Symptomatic infants have higher nasopharyngeal SARS-CoV-2 viral loads but less severe disease than older children. *Clin. Infect. Dis.* ciaa608 (2020) 818

757 29 T. Heald-Sargent *et al.*, Age-related differences in nasopharyngeal severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) levels in patients with mild to moderate coronavirus disease 2019 (COVID-19). *JAMA Pediatr.*, 10.1001/jamapediatrics.2020.3651 (2020). 819

758 30 Y. J. Park *et al.*; COVID-19 National Emergency Response Center, Epidemiology and Case Management Team, Contact tracing during coronavirus disease outbreak, South Korea, 2020. *Emerg. Infect. Dis.*, 10.3201/eid2610.201315 (2020). 820

759 31 M. Gale Jr, Interference with virus infection. *J. Immunol.* **195**, 1909–1910 (2015). 821

760 32 C. G. K. Ziegler *et al.*; HCA Lung Biological Network. Electronic address: lung-network@humancellatlas.org; HCA Lung Biological Network, SARS-CoV-2 receptor ACE2 is an interferon-stimulated gene in human airway epithelial cells and is detected in specific cell subsets across tissues. *Cell* **181**, 1016–1035.e19 (2020). 822

761 33 S. P. Sajuthi *et al.*, Type 2 and interferon inflammation strongly regulate SARS-CoV-2 related gene expression in the airway epithelium. *bioRxiv*:2020.04.09.034454 (10 April 2020). 823

762 34 A. Grifoni *et al.*, A sequence homology and bioinformatic approach can predict candidate targets for immune responses to SARS-CoV-2. *Cell Host Microbe* **27**, 671–680.e2 (2020). 824

763 35 A. Grifoni *et al.*, Targets of T cell responses to SARS-CoV-2 coronavirus in humans with COVID-19 disease and unexposed individuals. *Cell* **181**, 1489–1501.e15 (2020). 825

764 36 J. Mateus *et al.*, Selective and cross-reactive SARS-CoV-2 T cell epitopes in unexposed humans. *Science* **37**, eabd3871 (2020). 826

765 37 J. Braun *et al.*, SARS-CoV-2-reactive T cells in healthy donors and patients with COVID-19. *Nature*, 10.1038/s41586-020-2598-9 (2020). 827

766 38 K. W. Ng *et al.*, Pre-existing and *de novo* humoral immunity to SARS-CoV-2 in humans. *bioRxiv*:10.1101/2020.05.14.095414 (15 May 2020). 828

767 39 G. G. Wolff, Influenza vaccination and respiratory virus interference among Department of Defense personnel during the 2017-2018 influenza season. *Vaccine* **38**, 350–354 (2020). 829

768 40 C. S. Benn, M. G. Netea, L. K. Selin, P. Aaby, A small jab—a big effect: Nonspecific immunomodulation by vaccines. *Trends Immunol.* **34**, 431–439 (2013). 830

769 41 C. Pawlowski *et al.*, Exploratory analysis of immunization records highlights decreased SARS-CoV-2 rates in individuals with recent non-COVID-19 vaccinations. *medRxiv*:10.1101/2020.07.27.20161976 (29 July 2020). 831

770 42 R. Franklin *et al.*, Homologous protein domains in SARS-CoV-2 and measles, mumps and rubella viruses: Preliminary evidence that MMR vaccine might provide protection against COVID-19. *medRxiv*:10.1101/2020.04.10.20053207 (10 April 2020). 832

771 43 J. E. Gold, L. P. Tilley, W. H. Baumgartl, MMR vaccine appears to confer strong protection from COVID-19: few deaths from SARS-CoV-2 in highly vaccinated populations. *Researchgate*:10.13140/RG.2.2.32128.25607 (10 May 2020). 833

772 44 L. Steinman, A brief history of T(H)17, the first major revision in the T(H)1/T(H)2 hypothesis of T cell-mediated tissue damage. *Nat. Med.* **13**, 139–145 (2007). 834

773 45 Y. Du *et al.*, Clinical features of 85 fatal cases of COVID-19 from Wuhan. A retrospective observational study. *Am. J. Respir. Crit. Care Med.* **201**, 1372–1379 (2020). 835

774 46 C. Gruber *et al.*, Mapping systemic inflammation and antibody responses in multisystem inflammatory syndrome in children (MIS-C). *medRxiv*:10.1101/2020.07.04.20142752 (6 July 2020). 836

775 47 P. L. Acosta, M. T. Caballero, F. P. Polack, Brief history and characterization of enhanced respiratory syncytial virus disease. *Clin. Vaccine Immunol.* **23**, 189–195 (2015). 837

776 48 C. M. Szablewski *et al.*, SARS-CoV-2 transmission and infection among attendees of an overnight camp—Georgia, June 2020. *MMWR Morb. Mortal. Wkly. Rep.* **69**, 1023–1025 (2020). 838

777 49 V. G. Jones *et al.*, COVID-19 and Kawasaki disease: Novel virus and novel case. *Hosp. Pediatr.* **10**, 537–540 (2020). 839

778 50 J. Henry, R. J. Smeyne, H. Jang, B. Miller, M. S. Okun, Parkinsonism and neurological manifestations of influenza throughout the 20th and 21st centuries. *Parkinsonism Relat. Disord.* **16**, 566–571 (2010). 840

779 51 M. B. Son, K. Friedman, Coronavirus disease 2019 (COVID-19): Multisystem inflammatory syndrome in children. *Up to Date* (2020). <https://www.uptodate.com/contents/coronavirus-disease-2019-covid-10-multisystem-inflammatory-syndrome-in-children>. Accessed ■■■■. 841

780 Q:21 52 C. Bryce *et al.*, Pathophysiology of SARS-CoV-2: targeting of endothelial cells renders a complex disease with thrombotic microangiopathy and aberrant immune response. The Mount Sinai COVID-19 autopsy experience. *medRxiv*:10.1101/2020.05.18.20099960 (22 May 2020) 842

781 53 M. H. Han *et al.*, Proteomic analysis of active multiple sclerosis lesions reveals therapeutic targets. *Nature* **451**, 1076–1081 (2008). 843

782 54 L. Steinman, Platelets provide a bounty of potential targets for therapy in multiple sclerosis. *Circ. Res.* **110**, 1157–1158 (2012). 844

783 55 S. Garg *et al.*, Hospitalization rates and characteristics of patients hospitalized with laboratory-confirmed coronavirus disease 2019—COVID-NET, 14 states, March 1–30, 2020. *MMWR Morb. Mortal. Wkly. Rep.* **69**, 458–464 (2020). 845

784 56 D. A. Schwartz, An analysis of 38 pregnant women with COVID-19, their newborn infants, and maternal-fetal transmission of SARS-CoV-2: Maternal coronavirus infections and pregnancy outcomes. *Arch. Pathol. Lab Med.* **144**, 799–805 (2020). 846

785 57 J. A. W. Baidal *et al.*, Zooming towards a telehealth solution for vulnerable children with obesity during coronavirus disease 2019. *Obesity (Silver Spring)* **33**, 1184–1186 (2020). 847

786 848

787 849

788 850

789 851

790 852

791 853

792 854

793 855

794 856

795 857

796 858

797 859

798 860

799 861

800 862

801 863

802 864

803 865

804 866

805 867

806 868

# AUTHOR QUERIES

## AUTHOR PLEASE ANSWER ALL QUERIES

1

- Q: 1\_Please review 1) the author affiliation and footnote symbols, 2) the order of the author names, and 3) the spelling of all author names, initials, and affiliations and confirm that they are correct as set.
- Q: 2\_Please review the author contribution footnote carefully. Ensure that the information is correct and that the correct author initials are listed. Note that the order of author initials matches the order of the author line per journal style. You may add contributions to the list in the footnote; however, funding may not be an author's only contribution to the work
- Q: 3\_Please note that the spelling of the following author name in the manuscript differs from the spelling provided in the article metadata: Peggy Pui-Kay Ho. The spelling provided in the manuscript has been retained; please confirm.
- Q: 4\_Please review your open access and license selection and confirm that it is correct.
- Q: 5\_Certain compound terms are hyphenated when used as adjectives and unhyphenated when used as nouns. This style has been applied consistently throughout where (and if) applicable.
- Q: 6\_PNAS articles should be accessible to a broad scientific audience. As such, in the Abstract and at first occurrence in text, SARS-CoV-2 has been defined as severe acute respiratory syndrome coronavirus 2.
- Q: 7\_Affiliations: Please confirm or correct postal code added to each affiliation.
- Q: 8\_Please cite Fig. 1 in text.
- Q: 9\_PNAS articles should be accessible to a broad scientific audience. As such, CDC has been defined as Centers for Disease Control and Prevention, snRNA-seq has been defined as snRNA sequencing, and IFN has been defined as interferon. Please write out BLAST if it is an abbreviation.
- Q: 10\_“three different age groups ranging from approximately age 30 gestational weeks, approximately age 3 y, and approximately age 30 y”: “ranging from” requires a range “from a to b”; please rewrite accordingly or change to the following if the following is meant: “three different age groups: approximately age 30 gestational weeks, approximately age 3 y, and approximately age 30 y.”
- Q: 11\_“first described in 1957 by Isaacs and Lindemann (31)”: Because Isaacs and Lindemann are not the authors of ref. 31, this citation has been removed; the citation of ref. 31 at the end of the sentence has been retained.
- Q: 12\_“in children could provide some protection to SARS-CoV-2”: Should this be “. . .protection from SARS-CoV-2” to denote that the children, rather than SARS, are being protected? “some degree of protection to COVID-19” may similarly need to be changed.
- Q: 13\_“all had IgG to these to the receptor-binding domain of these common cold viruses”: This wording is confusing; you may wish to rewrite for clarity.



# AUTHOR QUERIES

## AUTHOR PLEASE ANSWER ALL QUERIES

2

- Q: 14\_“detected in individuals, including children, who were not exposed to SARS-CoV2”: If “who were not exposed. . .” only applies to children, please remove the comma after “children.”
- Q: 15\_The definition of IL has been moved to the first occurrence of the term, and the resulting later text “is mediated by IL-4 and IL-5 and IL-13” has been changed to “is mediated by IL-4, IL-5, and IL-13”; if IL-4 and IL-5 need to be paired together, separately from IL-13, please rewrite.
- Q: 16\_“associated with allergic disease including asthma”: Should “disease” be changed to “diseases”?
- Q: 17\_“A connection between knowledge gained in studying children”: “between” needs to be “between a and b”; please rewrite.
- Q: 18\_Ref. 1: Please note this report was updated on 14 August 2020 and the title now appears to be different from the version accessed on 11 August 2020.
- Q: 19\_Ref. 1: Please provide URL. Also, please note it appears this report was updated on 14 August 2020.
- Q: 20\_Refs. 6, 7, 25, 29, 30, and 37: Please provide volume and page numbers if article has been published.
- Q: 21\_Ref. 51: Please confirm reference as revised and provide date of access.
- 
-